

RESEARCH ARTICLE

Open Access Journal



Comparison of Nineteen Chemical Element Contents in Normal Thyroid and Thyroid with Riedel's Struma

Dr. Vladimir Zaichick PhD, DSc, FRSC, C.Chem*

¹Department of Radionuclide Diagnostics, Medical Radiological Research Center, Obninsk, Russia



Abstract

Background : Role of chemical elements (ChE) in etiology and pathogenesis of Riedel's disease (RD) is unclear.

Objectives: The aim of this exploratory study was to assess whether there were significant changes in thyroid tissue levels of nineteen ChE (Ag, Br, Ca, Cl, Co, Cr, Fe, Hg, I, K, Mg, Mn, Na, Rb, Sb, Sc, Se, Sr, and Zn) are present in the fibrotic transformed thyroid.

Methods : Nineteen ChE of thyroid tissue were determined in 6 patients with RD. The control group included thyroid tissue samples from 105 healthy individuals. Measurements were conducted using combination of non-destructive methods such as energy dispersive X-ray fluorescent analysis and instrumental neutron activation analysis.

Results : Reduced mean values of Ca and I content in 6.3 and 6.7 times, respectively, while elevated level of Br and Rb in 5.9 and 1.9 times, respectively, were found in thyroid with RD in comparison with normal level.

Conclusions : There are considerable changes in some ChE contents in tissue of thyroid with RD. Thus, it is reasonable to assume that the levels of these ChE in affected thyroid tissue can be used as RD markers. However, this topic needs additional studies.

Keywords: Riedel's disease, Intact thyroid, Chemical elements, Energy dispersive X-ray fluorescent analysis, Neutron activation analysis

Copyright : © 2021 The Authors. Published by Medical Editor and Educational Research Publishers Ltd. This is an open access article under the CC BY-NC-ND license

(<https://creativecommons.org/licenses/by-nc-nd/4.0/>).

1 | INTRODUCTION

Riedel's struma, also called Riedel's disease and Riedel's thyroiditis. is a peculiarly hard,

infiltrative lesion (nodule) of the thyroid gland (1). Riedel's disease (RD) is a rare form of chronic thyroiditis of unknown etiology associated with global or partial fibrosis of the thyroid gland, de-

COMPARISON OF NINETEEN CHEMICAL ELEMENT CONTENTS IN NORMAL THYROID AND THYROID WITH RIEDEL'S STRUMA

struction of the thyroid follicle architecture, obliterative phlebitis, and a mixed infiltrate of lymphocytes, eosinophils, and plasma cells (1, 2). Clinical differentiation between RD, Hashimoto's disease, and other thyroid benign and malignant nodules is often difficult (2, 3). We hypothesized that disbalance of trace elements (TE) contents in thyroid tissue may play a significant role in etiology and pathogenesis of RD. Furthermore, specific levels of TE contents in fibrotic transformed thyroid tissue may be used as RD biomarkers.

For over the 20th century, there was the governing opinion that all thyroid nodules (TN), including RD, are the straightforward sequel of iodine (I) deficiency. Though, it was found that TN is a frequent disease even in those countries and regions where the inhabitants are never exposed to I shortage (4). Moreover, it was shown that iodine excess has severe effects on human health and is associated with the development of thyroidal disfunctions and autoimmunity, nodular and diffuse goiter, benign and malignant tumors of gland (5–8). It was also demonstrated that besides the iodine deficiency and excess many other dietary, environmental, and occupational factors are associated with the TN incidence (9–11). Among them, a disruption of evolutionary stable input of many chemical elements (ChE) in the human body after the industrial revolution plays a significant role in the etiology of thyroidal disorders (12).

In addition to I, many other ChE is involved in essential physiological functions (13). Crucial or toxic (goitrogenic, mutagenic, carcinogenic) properties of ChE depend on tissue-specific need or tolerance, respectively (13). Deficiency, overload, or an imbalance of the ChE may result in cellular dysfunction, degeneration, death, benign or malignant transformation (13–15).

Supplementary information The online version of this article (<https://doi.org/10.52845/JMRHS/2021-4-11-3>) contains supplementary material, which is available to authorized users.

Corresponding Author: *Dr. Vladimir Zaichick PhD, DSc, FRSC, C.Chem*
Medical Radiological Research Centre Korolyev St.4, Obninsk 249036, Kaluga Region, Russia

In our earlier studies, the complex of in vivo and in vitro nuclear analytical and related methods was developed and used for the investigation of iodine and other ChE contents in the normal and pathological thyroid (16–22). Iodine level in the normal thyroid was scrutinized in relation to age, gender, and some non-thyroidal diseases (23, 24). Hereafter, variations of ChE content with age in the thyroid of males and females were studied, and age- and gender-dependence of some ChE was perceived (25–41). In addition, a significant difference between some ChE contents in normal and cancerous thyroid was demonstrated (42–47).

So far, the etiology and pathogenesis of RD has to be considered as multifactorial. The present study was performed to clarify the role of some ChE in the RD etiology. Having this in mind, our aim was to assess the silver (Ag), bromine (Br), calcium (Ca), chlorine (Cl), cobalt (Co), chromium (Cr), iron (Fe), mercury (Hg), I, potassium (K), magnesium (Mg), manganese (Mn), sodium (Na), rubidium (Rb), ammonium (Sb), scandium (Sc), selenium (Se), strontium (Sr), and zinc (Zn) contents in RD tissue using energy dispersive X-ray fluorescent analysis (EDXRF) combined with non-destructive instrumental neutron activation analysis with high resolution spectrometry of short-lived radionuclides (INAA-SLR) and long-lived radionuclides (INAA-LLR). A further aim was to compare the levels of these nineteen ChE in the RD transformed thyroid with those in intact (normal) thyroid (NT).

All studies were approved by the Ethical Committees of the MRRC. All the procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or with comparable ethical standards.

2 | MATERIAL AND METHODS

Samples

All patients with RD (n=6, 5 females and 1 male, mean age $M \pm SD$ was 39 ± 9 years, range 34–50) were hospitalized in the Head and Neck Department of the Medical Radiological Research Centre

(MRRC), Obninsk. Thick-needle puncture biopsy of suspicious lesion of the gland was performed for every persons, to allow morphological examination of affected thyroid tissue and to determine their TE contents. For all patients the diagnosis has been confirmed by clinical and morphological results obtained during studies of biopsy and resected materials. Histological conclusion for all thyroidal lesions was the RD.

Normal thyroids for the control group samples were drawn out at necropsy from 105 deceased (mean age 44 ± 21 years, range 2-87), who had died suddenly. The majority of deaths were owing to trauma. A histological examination in the control group was used to control the age norm conformity, also to confirm the absence of micro-nodules and latent cancer.

All tissue samples were divided into two parts using a titanium scalpel (48). One was used for morphological study, while the other was for ChE analysis. After the samples intended for ChE analysis were weighed, they were freeze-dried and homogenized (49).

Instrumentation and method

The content of Br, Fe, Rb, Sr, and Zn were determined by EDXRF. Details of the relevant facility for this method, source with ^{109}Cd radionuclide, methods of analysis and the results of quality control were presented in our earlier publications concerning the EDXRF of ChE contents in human thyroid and prostate tissue (25, 26, 50).

The content of Br, Ca, Cl, I, K, Mg, Mn, and Na were determined by INAA-SLR using a horizontal channel equipped with the pneumatic rabbit system of the WWR-c research nuclear reactor (Branch of Karpov Institute, Obninsk). Details of used neutron flux, nuclear reactions, radionuclides, gamma-energies, spectrometric unit, sample preparation and measurement were presented in our earlier publications concerning the INAA-SLR of ChE contents in human thyroid, scalp hair, and prostate (27, 28, 51–53)

In a few days after non-destructive INAA-SLR all thyroid samples were repacked and used for INAA-LLR. A vertical channel of the WWR-c research nuclear reactor (Branch of Karpov Institute, Obninsk). was applied to determine the content of Ag,

Co, Cr, Fe, Hg, Rb, Sb, Sc, Se, and Zn by INAA-LLR. Details of used neutron flux, nuclear reactions, radionuclides, gamma-energies, spectrometric unit, sample preparation and measurement were presented in our earlier publications concerning the INAA-LLR of ChE contents in human thyroid, scalp hair, and prostate (29, 30, 51, 54)

Standards and certified reference material

To determine contents of the ChE by comparison with a known standard, biological synthetic standards (BSS) prepared from phenol-formaldehyde resins were used (55). In addition to BSS, aliquots of commercial, chemically pure compounds were also used as standards. For each method ten certified reference material IAEA H-4 (animal muscle) and IAEA HH-1 (human hair) sub-samples were treated and analyzed in the same conditions that thyroid samples to estimate the precision and accuracy of results.

Statistical Analysis

A dedicated computer program for INAA mode optimization was used (56). All thyroid samples were prepared in duplicate, and mean values of ChE contents were used. Mean values of ChE contents were used in final calculation for the Br, Fe, Rb, and Zn mass fractions measured by two methods. Using Microsoft Office Excel, a summary of the statistics, including, arithmetic mean, standard deviation, standard error of mean, minimum and maximum values, median, percentiles with 0.025 and 0.975 levels was calculated for ChE contents. The difference in the results between two groups (NT and RD) was evaluated by the parametric Student's t-test and non-parametric Wilcoxon-Mann-Whitney U-test.

3 | RESULTS

resents certain statistical parameters of the Ag, Br, Ca, Cl, Co, Cr, Fe, Hg, I, K, Mg, Mn, Na, Rb, Sb, Sc, Se, Sr, and Zn mass fraction in normal thyroid and Riedel's struma.

Comparison of values obtained for Ag, Br, Ca, Cl, Co, Cr, Fe, Hg, I, K, Mg, Mn, Na, Rb, Sb, Sc, Se, Sr, and Zn contents in the NT samples with

COMPARISON OF NINETEEN CHEMICAL ELEMENT CONTENTS IN NORMAL THYROID AND THYROID WITH RIEDEL'S STRUMA

median of means reported by other researches (57–75) depicts in Table 2. A number of values for ChE mass fractions in literature were not expressed on a dry mass basis. However, we calculated these values using published data for water (75%) (76) and ash (4.16% on dry mass basis) (77) contents in thyroid of adults.

The ratios of means and the distinction between mean values of Ag, Br, Ca, Cl, Co, Cr, Fe, Hg, I, K, Mg, Mn, Na, Rb, Sb, Sc, Se, Sr, and Zn mass fractions in normal thyroid and Riedel's struma are presented in Table 3.

4 | DISCUSSION

Previously found good agreement of the Br, Ca, Cl, I, K, Mg, Mn, and Na contents analyzed by INAA-SLR with the certified data of CRM IAEA H-4 (18, 25–30, 50–54) indicates an acceptable accuracy of the results obtained in the study of ChE of the thyroid samples presented in Tables 1-3.

In a general sense values obtained for Br, Ca, Cl, Cr, Fe, Hg, I, K, Mg, Mn, Na, Rb, Sb, Sc, Se, and Zn contents in the normal human thyroid (Table 2) agree well with median of mean values reported by other researches (57–75). The obtained means for Ag and Co were almost one order of magnitude lower whereas mean for Sr was 7.46 times higher than median of previously reported means, but, nevertheless, inside the range of means (Table 2). Data cited in Table 2 for normal thyroid also includes samples obtained from patients who died from different non-endocrine diseases. In our previous study it was shown that some non-endocrine diseases can effect on ChE contents in thyroid (24). Moreover, in many studies the “normal” thyroid means a visually non-affected tissue adjacent to benign or malignant thyroidal nodules. However, there are no data on a comparison between the ChE contents in such kind of samples and those in thyroid of healthy persons, which permits to confirm their identity.

Table 1. Some statistical parameters of Ag, Br, Ca, Cl, Co, Cr, Fe, Hg, I, K, Mg, Mn, Na, Rb, Sb, Sc, Se, Sr, and Zn mass fraction (mg/kg, dry mass basis) in normal thyroid and Riedel's struma

Tissue	Element	Mean	SD	SEM	Min	Max	Median	P 0.025	P 0.975
Normal thyroid n=105	Ag	0.0151	0.0140	0.0016	0.0012	0.0800	0.0121	0.0017	0.0454
	Br	14.9	11.0	1.2	1.90	54.1	11.6	2.56	49.3
	Ca	1711	1022	109	414	6230	1458	460	3805
	Cl	3400	1452	174	1030	6000	3470	1244	5869
	Co	0.0399	0.0271	0.0030	0.0046	0.140	0.0327	0.0134	0.124
	Cr	0.539	0.272	0.032	0.130	1.30	0.477	0.158	1.08
	Fe	223	93	10	51.0	512	221	74.2	433
	Hg	0.0421	0.0358	0.0041	0.0065	0.180	0.0304	0.0091	0.150
	I	1841	1027	107	114	5061	1695	230	4232
	K	6071	2773	306	1740	14300	5477	2541	13285
	Mg	285	139	17	66.0	930	271	81.6	541
	Mn	1.35	0.54	0.07	0.510	4.18	1.32	0.537	2.23
	Na	6702	1764	178	3050	13453	6690	3855	10709
	Rb	8.16	4.55	0.49	1.66	29.4	7.37	3.08	19.3
	Sb	0.111	0.072	0.008	0.0047	0.308	0.103	0.0117	0.280
	Sc	0.0046	0.0038	0.0008	0.0002	0.0143	0.0042	0.00035	0.0131
	Se	2.32	1.29	0.14	0.439	5.80	2.01	0.775	5.65
	Sr	4.55	3.22	0.37	0.100	13.7	3.70	0.483	12.3
	Zn	105.1	40.1	4.3	7.10	221	104.9	39.2	186
Riedel's struma n=6	Ag	0.225	0.123	0.087	0.138	0.312	0.225	0.142	0.308
	Br	88.5	39.0	19.5	38.0	123	96.5	41.0	122
	Ca	279	238	168	111	447	279	119	439
	Cl	6252	3722	2149	3499	10487	4769	3563	10201
	Co	0.0588	0.0157	0.0111	0.0477	0.0699	0.0588	0.0483	0.0693
	Cr	1.14	1.05	0.74	0.396	1.88	1.14	0.433	1.84
	Fe	335	238	119	123	653	281	128	632
	Hg	0.529	0.088	0.063	0.466	0.591	0.529	0.469	0.588
	I	276	283	115	85	824	164	85.6	762
	K	12667	7652	4418	6612	21264	10111	6787	20706
	Mg	497	241	139	306	768	418	312	751
	Mn	2.05	1.30	0.75	0.57	3.92	2.57	0.67	3.00
	Na	6332	3642	2103	3732	10494	4769	3784	10208
	Rb	15.7	4.5	2.2	9.55	20.3	16.5	10.0	20.0
	Sb	0.0990	0.0198	0.0140	0.0850	0.113	0.0990	0.0857	0.112
	Sc	0.0150	0.0207	0.0150	0.00030	0.0296	0.0150	0.00103	0.0289
	Se	2.11	0.33	0.23	1.88	2.34	2.11	1.89	2.33
	Sr	10.4	10.6	5.3	1.09	23.2	8.57	1.18	22.6
	Zn	79.0	30.4	15.2	58.0	124	67.0	58.4	120

M – arithmetic mean, SD – standard deviation, SEM – standard error of mean, Min – minimum value, Max – maximum value, P 0.025 – percentile with 0.025 level, P 0.975 – percentile with 0.975 level.

Table 2 . Median, minimum and maximum value of means of twenty chemical element contents in the normal thyroid according to data from the literature in comparison with our results (mg/kg, dry mass basis)

Tissue	Element	Published data [Reference]			This work
		Median of means (n)*	Minimum of means M or M±SD, (n)**	Maximum of means M or M±SD, (n)**	
	Ag	0.21 (12)	0.000784 (16) [57]	1.20±1.24 (105) [58]	0.0151±0.0140
	Br	18.1 (11)	5.12 (44) [57]	284±44 (14) [59]	14.9±10.9
	Ca	1600 (17)	840±240 (10) [60]	3800±320 (29) [60]	1692±1022
	Cl	6800 (5)	804±80 (4) [61]	8000 (-) [62]	3400±1452
	Co	0.306 (25)	0.016 (66) [63]	70.4±40.8 (14) [59]	0.0399±0.0271
	Cr	0.69 (17)	0.088 (83) [64]	24.8±2.4 (4) [61]	0.539±0.272
	Cu	5.94 (61)	0.16 (83) [64]	220±22 (10) [61]	4.23±1.52
	Fe	252 (21)	56 (120) [65]	3360 (25) [66]	223±93
	Hg	0.08 (13)	0.0008±0.0002 (10) [60]	396±40 (4) [61]	0.0421±0.0358
	I	1888 (95)	159±8 (23) [67]	5772±2708 (50) [68]	1841±1027
	K	4400 (16)	46.4±4.8 (4) [61]	6090 (17) [69]	6071±2773
	Mg	390 (16)	3.5 (-) [70]	1520 (20) [71]	285±139
	Mn	1.62 (40)	0.076 (83) [64]	69.2±7.2 (4) [61]	1.35±0.58
	Na	8000 (9)	438 (-) [72]	10000±5000 (11) [73]	6702±1764
	Rb	7.8 (9)	=0.85 (29) [60]	294±191 (14) [59]	8.20±4.54
	Sb	0.15 (10)	0.040±0.003 (-) [72]	12.4 (-) [74]	0.111±0.072
	Sc	0.009 (4)	0.0018±0.0003 (17) [75]	0.014±0.005 (10) [60]	0.0046±0.0038
	Se	2.32 (21)	0.436 (40) [63]	756±680 (14) [59]	2.32±1.29
	Sr	0.61 (9)	0.055 (83) [64]	46.8±4.8 (4) [61]	4.55±3.22
	Zn	110 (56)	2.1 (-) [70]	820±204 (14) [59]	105±40

M – arithmetic mean, SD – standard deviation, (n)* – number of all references, (n)** – number of samples.

The data on ChE levels in RD tissue were not found in the literature.

The range of means of Ag, Br, Ca, Cl, Co, Cr, Fe, Hg, I, K, Mg, Mn, Na, Rb, Sb, Sc, Se, Sr, and Zn level reported in the literature for NT tissue vary widely (Table 2). This can be explained by a dependence of ChE content on many factors, including “normality” of thyroid samples (see above), the region of the thyroid, from which the sample was taken, age, gender, ethnicity, mass of the gland, and its functional activity. Not all these factors were strictly controlled in cited studies. However, in our opinion, the main reason for the inter-observer discrepancy can be attributed to the accuracy of the analytical techniques, sample preparation methods, and the inability to take standardized samples from affected tissues. It was insufficient quality control of results in these studies. In many scientific reports, tissue samples were ashed or dried at high temperature for many hours. In other cases, thyroid samples were treated with solvents (distilled water, ethanol, formalin etc). There is evidence that during ashing, drying and digestion at high temperature some quantities of certain ChE are lost as a result of this treatment. That concerns not only such volatile halogen as Br, but also other ChE investigated in the study (78, 79).

From Table 3, it is observed that in RD samples the mass fraction of Ca and I are approximately 6.3 and 6.7 times, respectively, lower, while Br and Rb content are 5.9 and 1.9 times, respectively, higher than in NT. Thus, if we accept the ChE contents in the NT group as a norm, we have to conclude that with a fibrotic transformation the Br, Ca, Rb, and I level in thyroid tissue notably changed.

Characteristically, elevated or reduced levels of ChE observed in affected tissues are discussed in terms of their potential role in the initiation and promotion of TN. In other words, using the low or high levels of the ChE in TN researchers try to determine the role of the deficiency or excess of each ChE in the TN etiology. In our opinion, abnormal levels of many ChE in TN, including RD, could be and cause, and also effect of thyroid tissue transformation. From the results of such kind studies, it is not always possible

to decide whether the measured decrease or increase in ChE level in pathologically altered tissue is the reason for alterations or vice versa. Nevertheless the differences between ChE levels in normal and affected thyroid tissue could be used as RD markers.

Table 3. Differences between mean values (M±SEM) of Ag, Br, Ca, Cl, Co, Cr, Fe, Hg, I, K, Mg, Mn, Na, Rb, Sb, Sc, Se, Sr, and Zn mass fraction (mg/kg, dry mass basis) in normal thyroid and Riedel's struma

Element	Thyroid tissue			Ratio	
	Normal thyroid	Riedel's struma	Student's t-test <i>p</i> ≤	U-test <i>p</i>	Riedel's struma to Normal thyroid
Ag	0.0151±0.0016	0.225±0.087	0.250	>0.05	14.9
Br	14.9±1.2	88.5±19.5	0.032	=0.01	5.94
Ca	1711±109	279±168	0.019	=0.01	0.16
Cl	3400±174	6252±2149	0.316	>0.05	1.84
Co	0.0399±0.0030	0.0588±0.0111	0.322	>0.05	1.47
Cr	0.539±0.032	1.14±0.74	0.567	>0.05	2.12
Fe	223±10	335±119	0.419	>0.05	1.50
Hg	0.0421±0.0041	0.529±0.063	0.080	>0.05	12.6
I	1841±107	276±115	0.00000002	=0.01	0.15
K	6071±306	12667±4418	0.274	>0.05	2.09
Mg	285±17	497±139	0.265	>0.05	1.74
Mn	1.35±0.07	2.05±0.75	0.451	>0.05	1.52
Na	6702±178	6332±2103	0.877	>0.05	0.94
Rb	8.16±0.49	15.7±2.2	0.041	=0.01	1.92
Sb	0.111±0.008	0.0990±0.0140	0.555	>0.05	0.89
Sc	0.0046±0.0008	0.0150±0.0150	0.607	>0.05	3.26
Se	2.32±0.14	2.11±0.23	0.531	>0.05	0.91
Sr	4.55±0.37	10.4±5.3	0.353	>0.05	2.29
Zn	105.1±4.3	79.0±15.2	0.184	>0.05	0.75

M – arithmetic mean, SEM – standard error of mean, Statistically significant values are in **bold**

5 | LIMITATIONS

This study has some limitations. Firstly, analytical techniques used in this study measure merely nineteen ChE (Ag, Br, Ca, Cl, Co, Cr, Fe, Hg, I, K, Mg, Mn, Na, Rb, Sb, Sc, Se, Sr, and Zn) mass fractions. Future studies should be aimed toward using other analytical methods such as inductively coupled plasma atomic emission spectrometry (ICP-AES) and inductively coupled plasma mass spectrometry (ICP-MS), which will elongate the list of ChE investigated in NT and RD. Secondly, the sample size of RD group was relatively small and prevented investigations of ChE contents in RD group using differentials like gender, thyroid functional activity, stage of disease, dietary habits of healthy persons and patients with RD. Lastly, the generalization of our outcomes may be bounded to the Russian population. Despite these limitations, this study provides

COMPARISON OF NINETEEN CHEMICAL ELEMENT CONTENTS IN NORMAL THYROID AND THYROID WITH RIEDEL'S STRUMA

evidence on fibrotic-specific tissue Br, Ca, Rb, and I level alteration and shows the necessity to continue ChE research of RD.

6 | CONCLUSION

In this work, ChE measurements were carried out in the tissue samples of normal thyroid and RD using three non-destructive instrumental analytical methods: EDXRF, INAA-SLR, and INAA-LLR. It was shown that the combination of these methods is an adequate analytical tool for the non-destructive determination of Ag, Br, Ca, Cl, Co, Cr, Fe, Hg, I, K, Mg, Mn, Na, Rb, Sb, Sc, Se, Sr, and Zn content in the tissue samples of human thyroid, including needle-biopsy samples. It was observed that in thyroid with RD content of Br and Rb significantly increased whereas the levels of Ca and I decreased in a comparison with the normal thyroid tissues. In our opinion, the increase in levels of Br and Rb, as well as the decrease in levels of Ca and I in RD transformed thyroid tissue might demonstrate an involvement of these ChE in etiology and pathogenesis of RD. It was supposed that the changes in levels Br, Ca, Rb, and I in thyroid tissue can be used as RD markers.

Declaration of Conflicting Interests

The author has not declared any conflict of interests.

7 | FUNDING

The author received no financial support for this study and for publication of this article.

ACKNOWLEDGEMENTS

The author is extremely grateful to Profs. Vtyurin BM and Medvedev VS, Medical Radiological Research Center, Obninsk, as well as to Dr. Choporov Yu, former Head of the Forensic Medicine Department of City Hospital, Obninsk, for supplying thyroid samples.

REFERENCES

1. Dahlgren M, Khosroshahi A, Nielsen P, Deshpande G, Stone V, H J. Riedel's thyroiditis and multifocal fibrosclerosis are part of the IgG4-related systemic disease spectrum. *Arthritis Care Res*;2010(9):1312–1318.
2. Junik R, Juraniec O, Pypkowski J, Krymer A, Marszałek A. A difficult diagnosis: a case report of combined Riedel's disease and fibrosing Hashimoto's thyroiditis. *Endokrynol Pol*. 2011;62(4):351–356.
3. Heufelder AE, Hay ID. Evidence for autoimmune mechanisms in the evolution of invasive fibrous thyroiditis (Riedel's struma). *Clin Investig*. 1994;72(10):788–793.
4. Derwahl M, Studer H. Multinodular goitre: 'much more to it than simply iodine deficiency. *Baillieres Best Pract Res Clin Endocrinol Metab*. 2000;14(4):577–600.
5. Zaichick V. Iodine excess and thyroid cancer. *J Trace Elem Exp Med*. 1998;11(4):508–509.
6. Zaichick V, Iljina T. Dietary iodine supplementation effect on the rat thyroid ¹³¹I blastomogenic action. *Die Bedeutung der Mengen- und Spurenelemente 18 Arbeitstangung Jena: Friedrich-Schiller-Universitat*. 1998;p. 294–306.
7. Kim S, Kwon YS, Kim JY, Hong KH, Park YK. Association between Iodine Nutrition Status and Thyroid Disease-Related Hormone in Korean Adults: Korean National Health and Nutrition Examination Survey VI. *Nutrients*. 2013;11(11):2757–2757.
8. Vargas-Uricoechea □, Pinzón-Fernández MV, Bastidas-Sánchez BE, Jojoa-Tobar E, Ramírez-Bejarano LE, Murillo-Palacios J. Iodine status in the colombian population and the impact of universal salt iodization: a double-edged sword? *J Nutr Metab*. 2019;p. 6239243–6239243.
9. Stojavljević A, Rovčanin B;. Available from: <https://doi.org/10.1007/s12403-021-00406-8>.

10. Fahim YA, Sharaf NE, Hasani IW, Ragab EA, Abdelhakim HK. Assessment of thyroid function and oxidative stress state in foundry workers exposed to lead. *J Health Pollut.* 2020;10(27).
11. Liu M, Song J, Jiang Y, Lin Y, Peng J, Liang H, et al. A case-control study on the association of mineral elements exposure and thyroid tumor and goiter. *Ecotoxicol Environ Saf.* 2021;208:111615–111615.
12. Zaichick V. Medical elementology as a new scientific discipline. *J Radioanal Nucl Chem.* 2006;269:303–309.
13. Moncayo R, Moncayo H. A post-publication analysis of the idealized upper reference value of 2.5 mIU/L for TSH: Time to support the thyroid axis with magnesium and iron especially in the setting of reproduction medicine. *BBA Clin.* 2017;7:115–119.
14. Beyersmann D, Hartwig A. Carcinogenic metal compounds: recent insight into molecular and cellular mechanisms. *Arch Toxicol.* 2008;82(8):493–512.
15. Martinez-Zamudio R, Ha HC. Environmental epigenetics in metal exposure. *Epigenetics.* 2011;6(7):820–827.
16. Zaichik VE, Yus R, Melnik AD, Cherkashin VI. Neutron-activation analysis in the study of the behavior of iodine in the organism. *Med Radiol (Mosk).* 1970;15(1):33–36.
17. Zaichik VE, Matveenko EG, Vtiurin BM, Medvedev VS. Intrathyroid iodine in the diagnosis of thyroid cancer. *Vopr Onkol.* 1982;28(3):18–24.
18. Zaichick V, Tsyb AF, Vtyurin BM. Trace elements and thyroid cancer. *Analyst.* 1995;120(3):817–821.
19. Vye Z, Yuya C. Determination of the natural level of human intra-thyroid iodine by instrumental neutron activation analysis. *J Radioanal Nucl Chem.* 1996;207(1):153–161.
20. Zaichick V. In vivo and in vitro application of energy-dispersive XRF in clinical investigations: experience and the future. *J Trace Elem Exp Med.* 1998;11(4):509–510.
21. Zaichick V, Zaichick S. Energy-dispersive X-ray fluorescence of iodine in thyroid puncture biopsy specimens. *J Trace Microprobe Tech.* 1999;17(2):219–232.
22. Zaichick V. Relevance of, and potentiality for in vivo intrathyroidal iodine determination. *Ann N Y Acad Sci.* 2000;904:630–632.
23. Zaichick V, Zaichick S. Normal human intrathyroidal iodine. *Sci Total Environ.* 1997;206(1):39–56.
24. Zaichick V. Human intrathyroidal iodine in health and non-thyroidal disease. *New aspects of trace element research.* 1999;p. 114–119.
25. Zaichick V, Zaichick S. Age-related changes of some trace element contents in intact thyroid of females investigated by energy dispersive X-ray fluorescent analysis. *Trends Geriatr Healthc;*2017(1):31–38.
26. Zaichick V, Zaichick S. Age-related changes of some trace element contents in intact thyroid of males investigated by energy dispersive X-ray fluorescent analysis. *MOJ Gerontol Ger.* 2017;1(5):28–28.
27. Zaichick V, Zaichick S. Age-related changes of Br, Ca, Cl, I, K, Mg, Mn, and Na contents in intact thyroid of females investigated by neutron activation analysis. *Curr Updates Aging.* 2017;1:5–6.
28. Zaichick V, Zaichick S. Age-related changes of Br, Ca, Cl, I, K, Mg, Mn, and Na contents in intact thyroid of males investigated by neutron activation analysis. *J Aging Age Relat Dis.* 2017;1(1):1002–1002.
29. Zaichick V, Zaichick S, Age, Ag, Co, Cr, et al. Se, and Zn contents in intact thyroid of females investigated by neutron activation analysis. *J Gerontol Geriatr Med.* 2017;3:15–15.

COMPARISON OF NINETEEN CHEMICAL ELEMENT CONTENTS IN NORMAL THYROID AND THYROID WITH RIEDEL'S STRUMA

30. Zaichick V, Zaichick S, Age, Ag, Co, Cr, et al. Se, and Zn contents in intact thyroid of males investigated by neutron activation analysis. *Curr Trends Biomedical Eng Biosci*. 2017;4(4):555644–555644.
31. Zaichick V, Zaichick S. Effect of age on chemical element contents in female thyroid investigated by some nuclear analytical methods. *MicroMedicine*. 2018;6(1):47–61.
32. Zaichick V, Zaichick S. Neutron activation and X-ray fluorescent analysis in study of association between age and chemical element contents in thyroid of males. *Op Acc J Bio Eng Bio Sci*. 2018;2(4):202–212.
33. Zaichick V, Zaichick S. Variation with age of chemical element contents in females' thyroids investigated by neutron activation analysis and inductively coupled plasma atomic emission spectrometry. *J Biochem Analyt Stud*. 2018;3(1):1–10.
34. Zaichick V, Zaichick S. Association between age and twenty chemical element contents in intact thyroid of males. *SM Gerontol Geriatr Res*. 2018;2(1):1014–1014.
35. Zaichick V, Zaichick S. Associations between age and 50 trace element contents and relationships in intact thyroid of males. *Aging Clin Exp Res*. 2018;30(9):1059–1070.
36. Zaichick V, Zaichick S. Possible role of inadequate quantities of intra-thyroidal bromine, rubidium and zinc in the etiology of female subclinical hypothyroidism. *EC Gynaecology*. 2018;7(3):107–115.
37. Zaichick V, Zaichick S. Possible role of inadequate quantities of intra-thyroidal bromine, calcium and magnesium in the etiology of female subclinical hypothyroidism. *Int Gyn and Women's. Health*. 2018;1(3).
38. Zaichick V, Zaichick S. Possible role of inadequate quantities of intra-thyroidal cobalt, rubidium and zinc in the etiology of female subclinical hypothyroidism. *Womens Health Sci J*. 2018;2(1):108–108.
39. Zaichick V, Zaichick S. Association between female subclinical hypothyroidism and inadequate quantities of some intra-thyroidal chemical elements investigated by X-ray fluorescence and neutron activation analysis. *Gynaecology and Perinatology*. 2018;2(4):340–355.
40. Zaichick V, Zaichick S. Investigation of association between the high risk of female subclinical hypothyroidism and inadequate quantities of twenty intra-thyroidal chemical elements. *Clin Res: Gynecol Obstet*. 2018;1(1):1–18.
41. Zaichick V, Zaichick S. Investigation of association between the high risk of female subclinical hypothyroidism and inadequate quantities of intra-thyroidal trace elements using neutron activation and inductively coupled plasma mass spectrometry. *Acta Scientific Medical Sciences*. 2018;2(9):23–37.
42. Zaichick V, Zaichick S. Trace element contents in thyroid cancer investigated by energy dispersive X-ray fluorescent analysis. *American Journal of Cancer Research and Reviews*. 2018;2:5–5.
43. Zaichick V, Zaichick S. Trace element contents in thyroid cancer investigated by instrumental neutron activation analysis. *J Oncol Res*. 2018;2(1):1–13.
44. Zaichick V, Zaichick S. Variation in selected chemical element contents associated with malignant tumors of human thyroid gland. *Cancer Studies*. 2018;2(1).
45. Zaichick V, Zaichick S. Twenty chemical element contents in normal and cancerous thyroid. *Int J Hematol Blo Dis*. 2018;3(2):1–13.
46. Zaichick V, Zaichick S. Levels of chemical element contents in thyroid as potential biomarkers for cancer diagnosis (a preliminary study). *J Cancer Metastasis Treat*. 2018;4:60–60.

47. Zaichick V, Zaichick S. Fifty trace element contents in normal and cancerous thyroid. *Acta Scientific Cancer Biology*. 2018;2(8):21–38.
48. Zaichick V, Zaichick S. Instrumental effect on the contamination of biomedical samples in the course of sampling. *The Journal of Analytical Chemistry*. 1996;51(12):1200–1205.
49. Zaichick V, Zaichick S. A search for losses of chemical elements during freeze-drying of biological materials. *J Radioanal Nucl Chem*. 1997;218(2):249–253.
50. Zaichick S, Zaichick V. The Br, Fe, Rb, Sr, and Zn contents and interrelation in intact and morphologic normal prostate tissue of adult men investigated by energy-dispersive X-ray fluorescent analysis. *X-Ray Spectrom*. 2011;40(6):464–469.
51. Zaichick S, Zaichick V. The effect of age and gender on 37 chemical element contents in scalp hair of healthy humans. *Biol Trace Elem Res*. 2010;134(1):41–54.
52. Zaichick S, Zaichick V. INAA application in the age dynamics assessment of Br, Ca, Cl, K, Mg, Mn, and Na content in the normal human prostate. *J Radioanal Nucl Chem*. 2011;288(1):197–202.
53. Zaichick V, Zaichick S. The effect of age on Br, Ca, Cl, K, Mg, Mn, and Na mass fraction in pediatric and young adult prostate glands investigated by neutron activation analysis. *J Appl Radiat Isot*. 2013;82:145–151.
54. Zaichick S, Zaichick V. The effect of age on Ag, Co, Cr, Fe, Hg, Sb, Sc, Se, and Zn contents in intact human prostate investigated by neutron activation analysis. *J Appl Radiat Isot*. 2011;69:827–833.
55. Zaichick V. Applications of synthetic reference materials in the medical Radiological Research Centre. *Fresenius J Anal Chem*. 1995;352:219–223.
56. Korelo AM, Zaichick V. Software to optimize the multielement INAA of medical and environmental samples. In: *Activation Analysis in Environment Protection*; 1993. p. 326–332.
57. Zhu H, Wang N, Zhang Y, Wu Q, Chen R, Gao J, et al. Element contents in organs and tissues of Chinese adult men. *Health Phys*. 2010;98(1):61–73.
58. Vlasova ZA. Dynamics of trace element contents in thyroid gland in connection with age and atherosclerosis. *Proceedings of the Leningrad Institute of Doctor Advanced Training*. 1969;80:135–144.
59. Salimi J, Moosavi K, Vatankhah S, Yaghoobi A. Investigation of heavy trace elements in neoplastic and non-neoplastic human thyroid tissue: A study by proton - induced X-ray emissions. *Int J Radiat Res*. 2004;1(4):211–216.
60. Boulyga SF, Zhuk IV, Lomonosova EM, Kanash NV, Bazhanova NN. Determination of microelements in thyroids of the inhabitants of Belarus by neutron activation analysis using the k0-method. *J Radioanal Nucl Chem*. 1997;222(1-2):11–14.
61. Reddy SB, Charles MJ, Kumar MR, Reddy BS, Ch A, Raju G, et al. Trace elemental analysis of adenoma and carcinoma thyroid by PIXE method. *Nuclear Instruments and Methods in Physics Research Section B: Beam Interactions with Materials and Atoms*. 2002;196(3-4):333–339.
62. Woodard HQ, White DR. The composition of body tissues. *Brit J Radiol*. 1986;708:1209–1218.
63. Stojavljević A, Rovčanin B, Krstić D, Borković-Mitić S, Paunović I, Diklić A, et al. Risk assessment of toxic and essential trace metals on the thyroid health at the tissue level: The significance of lead and selenium for colloid goiter disease. *Expo Health*. 2019;.
64. Reitblat MA, Kropachyev AM. Some trace elements in thyroid of the Perm Pricam'ya residents. *Proceedings of Perm Medical Institute*. 1967;78:157–164.

COMPARISON OF NINETEEN CHEMICAL ELEMENT CONTENTS IN NORMAL THYROID AND THYROID WITH RIEDEL'S STRUMA

65. Ataulachanov IA. Age changes in the content of manganese, cobalt, copper, zinc and iron in the endocrine glands of women. *Probl Endocrinol (Mosk)*. 1969;15(2):98–102.
66. Kamenev VF. About trace element contents in thyroid of adults. In: *Trace Elements in Agriculture and Medicine*; 1963. p. 12–16.
67. Neimark II, Timoschnikov VM. Development of carcinoma of the thyroid gland in person residing in the focus of goiter endemic. *Problemy Endocrinologii*. 1978;24(3):28–32.
68. Zabala J, Carrion N, Murillo M, Quintana M, Chirinos J, Seijas N, et al. Determination of normal human intrathyroidal iodine in Caracas population. *J Trace Elem Med Biol*. 2009;23(1):9–14.
69. Forssen A. Inorganic elements in the human body. *Ann Med Exp Biol Fenn*. 1972;50(3):99–162.
70. Kortev AI, Dontsov GI, Lyascheva AP. Bioelements and a human pathology. Sverdlovsk, Russia; 1972.
71. Li AA. Level of some macro- and trace element contents in blood and thyroid of patients with endemic goiter in Kalinin region. Kalinin, Russia; 1973.
72. Boulyga SF, Becker JS, Malenchenko AF, Dietze HJ. Application of ICP-MS for multi-element analysis in small sample amounts of pathological thyroid tissue. *Microchimica Acta*. 2000;134(3-4):215–222.
73. Soman, Joseph KT, Raut SJ, Mulay CD, Parameshwaran M, Panday VK. Studies of major and trace element content in human tissues. *Health Phys*. 1970;19(5):641–656.
74. Zakutinsky DK, Parfyenov YuD, Selivanova LN. *Handbook of the toxicology of radioactive isotopes*. Moscow: State Publishing House of Medical Literature; 1962.
75. Kvicala J, Havelka J, Nemeč J, Zeman V. Selenium and rubidium changes in subjects with pathologically altered thyroid. *Biol Trace Elem Res*. 1992;32:253–258.
76. Katoh Y, Sato T, Yamamoto Y. Determination of multielement concentrations in normal human organs from the Japanese. *Biol Trace Elem Res*. 2002;90(1-3):57–70.
77. Schroeder HA, Tipton IH, Nason AP. Trace metals in man: strontium and barium. *J Chron Dis*. 1972;25(9):491–517.
78. Zaichick V. Sampling, sample storage and preparation of biomaterials for INAA in clinical medicine, occupational and environmental health. In: *Harmonization of Health-Related Environmental Measurements Using Nuclear and Isotopic Techniques*. Vienna: IAEA; 1997. p. 123-133
79. Zaichick V. Losses of chemical elements in biological samples under the dry aching process. *Trace Elements in Medicine*. 2004;5(3):17–22.

How to cite this article: Zaichick V.. **Comparison of Nineteen Chemical Element Contents in Normal Thyroid and Thyroid with Riedel's Struma**. *Journal of Medical Research and Health Sciences*. 2021;1529–1538. [ht tps://doi.org/10.52845/JMRHS/2021-4-11-3](https://doi.org/10.52845/JMRHS/2021-4-11-3)