



Angina Bullosa Hemorrhagica in Oral Cavity: A Case Report

*Abdulaziz A Alwakeel¹, Abdulazez A Aleid², Mohammed Rahoof³, Khalid Alenezi⁴, Anwar Alanazi⁵

¹Oral medicine specialist, oral medicine and pathology department. Tabuk specialist dental center, Tabuk health cluster, Tabuk city, Saudia Arabia

²Oral and maxillofacial surgeon, Oral and maxillofacial surgery department. Tabuk specialist dental center, Tabuk health cluster, Tabuk city, Saudia Arabia

³Pediatric dentist, pediatric dentistry department. Tabuk specialist dental center, Tabuk health cluster, Tabuk city, Saudia Arabia

⁴Postgrad endodontic resident, endodontic department., Tabuk specialist dental center, Tabuk health cluster, Tabuk city, Saudia Arabia

⁵Dental resident, Public health department. Tabuk specialist dental center, Tabuk health cluster, Tabuk city, Saudia Arabia

This study was done in Oral medicine and pathology department, Tabuk specialist dental center, Tabuk health cluster, Tabuk city, Saudia Arabia



Abstract

Angina bullosa hemorrhagica (ABH) was primarily introduced by Badham in 1967; it is a rare, idiopathic condition in the oral cavity, which was described as a bulla filled with blood. The present Case shows A 33-year-old healthy male patient presented to the dental clinic with an oral lesion in the palatal area that had been present for 4 days. The history revealed that the lesion appeared after eating hard and spicy food. This was the patient's first episode of such swelling. The patient had no history of chronic bleeding disorders and was not on any regular medications. Clinical examination revealed a vesicobullous lesion on the left side of the soft palate, which was dark blue in color, filled with blood, and measured 1.5 x 1 cm in diameter. The patient was treated with prednisolone 5 mg once a day for 3 days as a mouthwash, along with chlorhexidine 0.12% twice a day for 5 days. A follow-up appointment after 2 weeks showed complete healing of the lesion

Keywords: Oral bullosa hemorrhagica, soft palate hemorrhagica, soft palate lesion

Copyright: © 2021 The Authors. Published By Medical Editor And Educational Research Publishers Ltd. This Is An Open Access Article Under The Cc By-Nc-Nd License (<https://creativecommons.org/licenses/by-nc-nd/4.0/>).

Corresponding Author: Abdulaziz A Alwakeel

Angina Bullosa Hemorrhagica in Oral Cavity: A Case Report

Introduction:

Angina bullosa hemorrhagica (ABH) was primarily introduced by Badham in 1967; it is a rare, idiopathic condition in the oral cavity, which was described as a bulla filled with blood.[1]

This lesion characterized as a dark red-purple colored bulla, which is a filled bulla and surrounded by a red margin with approximate measures between 1 to 3cm. [2]It is commonly seen during the fifth decade of life .[3]

The lesion is caused by bleeding in the superficial capillaries of blood vessels due to traumatic injury to the epithelial-connective-tissue junction, thus resulting in subepithelial hemorrhagic bulla formation.[4] Until now, the etiopathogenesis of this lesion has not been clear.[2]

It was also thought that it starts spontaneously or occurs with a traumatic factor primarily due to hot or hard food.[5]

On other hand , the most common affected site in the oral cavity is the soft palate .[6]

No biopsy is required for most of ABH cases .[7]

Many differential diagnoses can be considered, such as thrombocytopenia and autoimmune diseases like pemphigoid and epidermolysis bullosa.[8]

The aim of this paper is to present a case of ABH in a healthy patient, caused by traumatic injury, and treated with a combination of chlorhexidine and corticosteroid as a mouthwash.

Case Presentation:

A 33-year-old healthy male patient presented to the dental clinic with an oral lesion in the palatal area that had been present for 4 days. The history revealed that the lesion appeared after eating hard and spicy food. This was the patient's first episode of such swelling. The patient had no history of chronic bleeding disorders and was not on any regular medications. There was no family history of chronic diseases. Clinical examination revealed a vesicobullous lesion on the left side of the soft palate, which was dark blue in color, filled with blood, and measured 1.5 x 1 cm in diameter, as shown in Figure 1.



Figure 1: Clinical photograph shows dark blue colored bulla lesion on the left side of the soft palate

The patient was treated with prednisolone 5 mg once a day for 3 days as a mouthwash, along with chlorhexidine 0.12% twice a day for 5 days. A

follow-up appointment after 2 weeks showed complete healing of the lesion, as shown in **Figure2**

Angina Bullosa Hemorrhagica in Oral Cavity: A Case Report




Figure 2 : Complete healing at the lesion site after 2 weeks of treatment

Written and verbal consent were obtained from the patient to publish his clinical and histopathological images for scientific purposes. Additionally, IRB approval was granted by the Tabuk Institutional Review Board under protocol No. TU-077/024/252 for this case report.

Discussion

ABH tends to occur in adults between the fifth and the seventh decades of life, with a mean age of 55.4 years. With equal distribution between males and females. [9]

The lesions usually healed in short time without scar. [6] These lesions need quick intervention as it can cause airway obstruction. [9]

ABH may have an association with systemic diseases like diabetes mellitus and hypertension. [10]

Usage of inhaled corticosteroids for long time can cause in impair collagen formation and later atrophy of the epithelial tissue. [11]

Collagen and elastic fibers disorder in the oral cavity results in less anchorage of blood vessels, which can cause hemorrhagic lesions after trauma. [12]

Analgesic medication with antibiotic mouthwash, such as (chlorhexidine 0.12-0.2%) can be used. Large intact lesions, especially on the soft palate, should be incised and drained to avoid a possible obstruction of the upper aerodigestive tract. [13]

Conclusion : In the present case, this lesion may be associated with a systemic disease, particularly of a dermatological nature. A thorough systemic evaluation is indicated. Topical treatments, such as chlorhexidine 0.12% along with corticosteroids as a mouthwash, can be effective in treating this type of oral lesion.

References:

1. Badham NJ. Blood blisters and the oesophageal cast. *J Laryngol Otol* 1967; 81(7): 791–803.
2. Beguerie JR and Gonzalez S. Angina bullosa hemorrhagica: report of 11 cases. *Dermatol Rep* 2014; 6(1): 5282.
3. Horie N, Kawano R, Inaba J, et al. Angina bullosa hemorrhagica of the soft palate: a clinical study of 16 cases. *J Oral Sci* 2008; 50 (1): 33–36
4. Martini MZ, Lemos CA and Jr Shinohara EH. Angina bullosa hemorrhagica: report of 4 cases. *Minerva Stomatol* 2010;59(3):139–142
5. Ordioni U, Hadj Saïd M, Thiery G, et al. Angina bullosa haemorrhagica: a systematic review and proposal for diagnostic criteria. *Int J Oral Maxillofac Surg* 2019; 48(1): 28–39
6. Yamamoto K, Fujimoto M, Inoue M, et al. Angina bullosa hemorrhagica of the soft

Angina Bullosa Hemorrhagica in Oral Cavity: A Case Report

- palate: report of 11 cases and literature review. *J Oral Maxillofac Surg* 2006;64(9):1433–1436.
7. Giuliani M, Favia GF, Lajolo C, et al. Angina bullosa haemorrhagica: presentation of eight new cases and a review of the literature. *Oral Dis* 2002; 8(1): 54–58.
 8. Mukhopadhyaya I and Rath CS. Angina bullosa hemorrhagica-an uncommon oral lesion. *Indian J Dent Adv* 2018; 10(1): 60–63
 9. Cinar SL, Kartal D, Canöz Ö, Borlu M, Ferahbas A. Case Report: A rare cause of oral bullae: Angina bullosa hemorrhagica. *F1000 Res.* 2017 Nov 8;(6):1974..
 10. Grinspan D, Abulafia J, Lanfranchi H. Angina bullosa hemorrhagica. *Int J Dermatol.* 1999 Jul;38(7):525-8.
 11. Antoni-Bach N, Couilliet D, Garnier J, Tortel MC, Grange F, Guillaume JC. Cas pour diagnostic. Stomatite bulleuse hémorragique bénigne (SBHB) [Case for diagnosis. Benign hemorrhagic bullous stomatitis]. *Ann Dermatol Venereol.* 1999 Jun-Jul;126(6-7):525-6
 12. Higgins EM, du Vivier AW. Angina bullosa haemorrhagica--a possible relation to steroid inhalers. *Clin Exp Dermatol.* 1991 Jul;16(4):244-6.
 13. Pahl C, Yarrow S, Steventon N, Saeed NR, Dyar O. Angina bullosa haemorrhagica presenting as acute upper airway obstruction. *Br J Anaesth.* 2004 Feb;92(2):283-6.

How to Cite: Alwakeel, A., Aleid, A. A., Rahoof, M., Alenezi, K., & Alanazi, A. (2024). Angina bullosa hemorrhagica in oral cavity: A case report. *Jour Med Resh and Health Sci*, 7(12),3287–3290. <https://doi.org/10.52845/JMRHS/2024-7-12-4>